

Milk metabolites are non-invasive biomarkers for nutritional and metabolic disorders of dairy herds?

Pinheiro, C.^{1*}, Domingues, I.^{2,3**}, Vaz, P.³, Moreira, R.³ & Infante, P.⁴

*ccp@uevora.pt; **inesdomingues@eabl.pt

What?

The interpretation of milk metabolites from milk recording can be indicative of nutritional and metabolic disorders?

Who?

University of Évora
and EABL

Why?

Milk Recording is done monthly in dairy farmers, is this a management tool?



Where?

27 herds in South
of Portugal

How?

Individual milk samples collected
monthly from January to March
2017

Background



J. Dairy Sci. 99:9263–9270
<http://dx.doi.org/10.3168/jds.2016-11128>
© American Dairy Science Association®, 2016.

Prevalence of elevated milk β -hydroxybutyrate concentrations in Holstein cows measured by Fourier-transform infrared analysis in Dairy Herd Improvement milk samples and association with milk yield and components

D. E. Santschi,¹ R. Lacroix, J. Durocher, M. Duplessis, R. Valacta, 555, boul. Des Anciens-Combattants, Ste-Anne-de-Bellevue, Québec



Revista Brasileira de Zootecnia

© 2014 Sociedade Brasileira de Zootecnia
ISSN 1806-9290
www.sbz.org.br

R. Bras. Zootec., 43(9):479-484, 2014

Lactose and milk urea nitrogen: fluctuations during lactation in Holstein cows

Andrés F. Henao-Velásquez¹, Oscar David Múnera-Bedoya¹, Ana Cristina Herrera¹, Jorge H. Agudelo-Trujillo², Mario Fernando Cerón-Muñoz^{1,2}

¹Research Group, Facultad de Ciencias Agrarias, Universidad de Antioquia, Medellín, Colombia.
²Ciencias Agrarias, Universidad de Antioquia, Medellín, Colombia.

THE IMPORTANCE OF MONITORING CHANGES IN MILK FAT TO MILK PROTEIN RATIO IN HOLSTEIN COWS DURING LACTATION

VÝZNAM SLEDOVÁNÍ ZMĚN POMĚRU TUK/BÍLKOVINA V MLÉČE HOLŠTÝNSKÝCH DOJNIC

Vladimír ČEJNA*, Gustav CHLÁDEK

Department of Animal Breeding, Mendel University of Agriculture and Forestry in Brno, Zemědělská 1, 613 00 Brno, Czech Republic, Tel.: +420 545 133 220, e-mail: xcejnav@seznam.cz

Manuscript received: October 3, 2005; Reviewed: October 10, 2005; Accepted for publication: October 10, 2005



ELSEVIER

Preventive Veterinary Medicine 59 (2003) 83–93

www.elsevier.com/locate/prevetmed

PREVENTIVE
VETERINARY
MEDICINE

The effect of non-nutritional factors on milk urea nitrogen levels in dairy cows in Prince Edward Island, Canada

P. Arunvipas*, I.R. Dohoo, J.A. VanLeeuwen, G.P. Keefe

Department of Health Management, Atlantic Veterinary College, University of Prince Edward Island (UPEI), 550 University Avenue, Charlottetown, PEI, Canada C1A 4P3

Received 28 February 2002; received in revised form 10 December 2002; accepted 26 January 2003




EAAP 2018

69th Annual Meeting of the European Federation of
Animal Science

Dubrovnik, Croatia, 27th to 31st August 2018

Background



Nutrient Imbalances - relationship between carbohydrates fermentability and protein degradability in the rumen can be diagnosed by **Milk Urea Nitrogen, Protein and Relation of Fat/Protein.**

Metabolic Imbalances - Negative Energy Balance, Hyperketonemia and Ketosis can be diagnosed by **β -hydroxybutyrate, Fat and Relation of Fat/Protein.**

Metabolic Imbalances - Acidosis can be diagnose by **Fat and Relation of Fat/Protein.**

Milk metabolites can be indicators of **Health and Welfare** of the cow

Background



Thresholds


Levels	Fat (%)	Protein (%)	Relation F/P	MUN (mg/kg)	BHB (mmol/L)
Low	< 3,20 ¹	< 3,00 ^{3,4,5}	< 1,00 ⁶	< 100	< 0,10 ⁸
Normal	3,20 – 4,50	3,00 – 3,20	1,00 – 1,40	100 – 300	0,10 – 0,20
High	> 4,50 ²	> 3,20 ^{3,4,5}	> 1,40 ^{1,7}	> 300	> 0,20 ^{9,10,11,12}

Material & Methods



- 27 farms
- 1 milk sample / month / cow
- January 2015 – March 2017
- 110461 individual milk samples - 9523 lactation

- Descriptive Statistics
- Boxplots
- GLMM
- Spearman's Correl.



Results



First 30 days of lactation:

7.7% of milk recording had BHB concentration over 0.2mmol/L.
These cows had high possibility of being with clinical ketosis.



44.8% of milk recording had the relation of F/P over 1.4.
49.3% of milk recording had milk fat over 4.5%.
These cows were probability mobilizing body fat.



11.9% of milk recording had the relation of F/P above 1.4.
21.6% of milk recording had milk protein above 3%.
These cows were ingesting a small proportion of protein in the diet comparing with the quantity of carbohydrates.

Conclusions



These non-invasive biomarkers can reflect **nutritional and metabolic disorders.**



The interrelation between them must be taken into account.
The thresholds of this milk metabolites to indicate health disorders are not consensual among the authors.



ccp@uevora.pt



inesdomingues@eabl.pt



EAAP 2018

69th Annual Meeting of the European Federation of
Animal Science

Dubrovnik, Croatia, 27th to 31st August 2018