Benefits and limits of farm animals to control herbage mass, pests and weeds in orchards: a review

- Gentiane Maillet (ACTA)
- Arnaud Dufils (INRAE)
- Frédérique Angevin (INRAE)
- Sonia Ramonteu (ACTA)
- Jean-Louis Peyraud (INRAE)
- <u>René Baumont (INRAE)</u>

















Background of the study

Grandes

- Exploring synergies between livestock and crop sectors is seen as a promising way towards a more sustainable and circular European agriculture (ATF and Plant ETP, 2019, 2020)
- Initiatives aimed at reconnecting livestock and crop production at different scales are emerging, but remain scattered, making it difficult to assess and disseminate them
- The four GIS on agricultural sectors led by INRAE (Avenir Elevages, Grandes Cultures, Fruits et PIClég), in collaboration with ACTA and the thematic network on mixed-crop farming (RMT SPICEE), have joined forces to carry out a project focusing on the reconnection between crops and livestock (<u>https://www.gis-avenir-elevages.org/Livestock-Farming-Future-Who-arewe/REVE-REconnexion-Vegetal-Elevage</u>)





RMT SPICEE

Context and objectives

- One of the levers of the agroecological transition is the diversification of systems, in particular by reintroducing animals into systems specialized in field crops or arboriculture
- In this study, we focus on the introduction of animals into fruit and vineyard plots
- One of the assumed benefits of introducing animals into the plots is the control of herbage mass and weeds and the management of pests through biological regulation
- But animals can also cause damages on soil and vegetation

→ Make a comparative synthesis of the benefits and limits of different animals (small and large herbivores, poultry, pigs) to control herbage mass, weeds and pests in vineyards and orchards



Methods : 66 documents collected

- 23 Scientific publications
- 18 Technical publications or deliverables from R&D projects
- 23 Testimonies in in professional journals and in social media
- 2 Interviews conducted with ongoing experimental projects

→ Geographical distribution : France (40), EU (9), Rest of the world (17)

















« Chickens in orchards », MIRAD project « Rabbits in orchards », LAPOESIE project





Photos : © INRAE - Sara Bosshardt



Methods : analysis of the documents

Construction of a matrix of the effects of animal species on orchards and vineyards:

- Ruminants (sheep, cows, goats)
- Poultry (chickens, geese, guinea fowl, ducks)
- Pigs
- Rabbits
- Equines (horses, donkeys)

- Herbage mass and grass cover
- Pests (rodents, insects and other invertebrates)
- Pathogens
- Auxiliary species
- Risks of damages to trees and soil

	Very positive effects (only benefits mentioned)
	Positive effects (benefits mentioned but not systematic)
	No effects
	Negative effects (disadvantages mentioned but solutions proposed)
	Very negative effects (disadvantages mentioned but no solutions proposed)
	Unknown effects
	No data available
(?)	Hypothesis to be confirmed

Numbers correspond to the number of documents









General overview



	- Grass consumption	37	7	11	14	9	(?) 5	(?) 1	(?) 2	10	5	1
	- Exploring the plot	37	7	11	14	9	(?) 5	1	(?) 2	10	5	1
Grass cover / soil	- Appearance of refusals	14		1	3	3	(?) 2	(?) 1	(?) 2		1	
fertility	Changes in plant diversity	3			2	2						
lerency	-Thinning out the leaves	2										
	- Improvement in soil fertility (looser, more aerated soil) by scraping or trampling	1			3							
	- Destruction of galleries by trampling	5		1						(?) 2		
	- Destroying galleries by excavating the ground									(?) 2		
and a	- Rodent consumption									(?) 2		
	- Rejuction in the vegetation cover that protects rodents from natural predation	2		1						(?) 1		
	- Reduction of natural predation by rodents through the use of fences restricting access to plots of land				1							
Pests	- Bio ogical regulation of plant pathogenic nematodes by other nematode species	1										
	- Corsumption of insects, larvae or molluscs			2	(?) 13	4	4		2	(?) 1	1	
	- Corsumption of infested fruit that has fallen to the ground	(?) 9		2	(?) 8	(?) 6	(?) 2	(?) 1	1	(?) 4		
	- Disturbance of insects by rubbing against trunks	2										
71.	- Disturbance to insect habitat and microclimate as a result of reduced grass cover				1	1	1					
х. 	- Appearance of insect pests living in animal dung			1								
	- Reappearance of auxiliary species (unknown mechanisms)	1										
Auxiliary	- Consumption of auxiliary species				(?) 2							
species 🔫	- Changes in auxiliary species due to changes in flora	(?) 1										
	-Reducing inoculum by trampling or eating dead leaves or fruit that has fallen to the ground	(?) 9			(?) 3	(?) 2				(?) 4	(?) 2	
Pathogens	- Reduction of inoculum by the degradation of dead leaves that have fallen to the ground thanks to urine (droppings in the case of poultry).	(?) 1			(?) 1	(?) 1				(?) 1	(?) 1	
	- Improving the microclimate by removing low branches	5										
	- Reduced risk of ascospore projection due to removal of low branches	2										
	- Improving the microclimate by removing the leaves from the vine shoots	1										
	 Root damage caused by "ploughing" the soil or scratching the surface 				(?) 3		1			7		
	- Degradation of the tree by barking or pecking	17	3	2	(?) 6	3	3	1	2		2	
	- Damage to low branches caused by consumption, rubbing or leaning against the tree	18	5	4	3	3	2	1	2			
	- Consumption of buds, leaves or fruit	20	5	4	(?) 7	3	2	1	2	2		1
	- Damage to the trunk by pressure or friction	4		2	(?) 4	2	1	1	1			
Rick of damages	- Damage to the tree by perching				2		1					
INISK OF GAILIAges	- Damage to vine shoots	1	1							1		
- AND	- Soil degradation through compaction or the creation of ruts	13	2	4	2	1	1			7		
	- Appearance of bare soil	2			(?) 4							
	- Damage to fruit	1			(?) 2		(?) 1					
	- Damage to young plants	5	1	1	3	2	2	1	1			
'	- Degradation of trees through local over-fertilisation	2										

Grass cover management

Sheep, cows, geese, chickens, rabbits, are interesting species for herbage management

- Sheep: a traditional and effective solution, provided that the risk of damage is kept under control
- Cows: animals traditionally used in high-stem orchards or coconut plantations
- Geese: herbivorous poultry that are easy to introduce
- Chickens or guinea fowl: a possible solution, but may result in bare soil
- Rabbits: an innovative way of managing grass cover in orchards

Goats and pigs: not recommended animals for managing grass cover because of the damage they cause to trees and soil respectively





Photo : © INRAF - Sara Bosshard













Benefits for pest and disease management

Rodent management

• Destruction of rodent galleries by trampling: the most frequently reported mechanism

Direct or indirect action by animals on insects or other invertebrates

• Some animals (poultry) are heavy consumers of larvae, insects or mollusks

The introduction of animals into orchards also plays a prophylactic role

- The consumption or trampling of dead leaves or fruit that has fallen to the ground helps to reduce scab inoculum
- Eating low branches or pruning them limits the spread of disease.



Major knowledge gaps 🐐 🖗 🌾 🖉

-Grass consumption 37 7 11 14 9 0,15 0,11 0,2 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 <t< th=""><th></th><th></th><th></th><th>÷</th><th></th><th></th><th></th><th></th><th></th><th></th><th></th><th></th><th></th></t<>				÷									
Grass cover / sol fertility - sploring the plot. 97 11 14 9 (P) 2 10 2 Grass cover / sol fertility - Sploring the plot. 33 12 3 0(P) 2 (P) 2 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10 10		- Grass consumption	37	7	11	14	9	(?) 5	(?) 1	(?) 2	10	5	1
Grass cover / soil - Appearance of refusals 14 1 3 3 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 2 (?) 1 (?) 2 (?) 2 (?) 1 1 (?) 2 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 </td <td></td> <td>- Exploring the plot</td> <td>37</td> <td>7</td> <td>11</td> <td>14</td> <td>9</td> <td>(?) 5</td> <td>1</td> <td>(?) 2</td> <td>10</td> <td>5</td> <td>1</td>		- Exploring the plot	37	7	11	14	9	(?) 5	1	(?) 2	10	5	1
fertility - Changes in plant diversity. 3 2 2 Importance Improvement in soil fertility (loser, more aerated soil) by scraping or trampling 1 3	Grass cover / soi	- Appearance of refusals	14		1	3	3	(?) 2	(?) 1	(?) 2		1	
Increment 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	fortility	- Changes in plant diversity	3			2	2						
Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 3 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or trampling 1 Improvement in solf ertility (losser, more aerated soil) by scraping or scratting scrapics 1 Improvement in solf ertility (losser, more aerated soil) by scraping 1 Improvement in solf ertility (losser, more aerated soil) by scraping 1 1	lerency	- Thinning out the leaves	2										
 Destruction of galleries by trampling. Destruction of galleries by excavating the ground Reduction in the vegetation cover that protects rodents from natural predation Reduction of natural predation by rodents through the use of fences restricting access to plots of land Reduction of inset est privation of plant pathogenic nematodes by other nematode species Consumption of insects, larvae or molluscs Consumption of insects of trubing against trunks Disturbance of insects by trubing against trunks Disturbance of insects by trubing against trunks Disturbance of insects by trubing against trunks Consumption of auxiliary species (unknown mechanisms) Consu		- Improvement in soil fertility (looser, more aerated soil) by scraping or trampling	1			3							
 Destroying galleries by excavating the ground Pests Biological regulation of plant pathogenic nematodes by other nematode species Consumption of lanet prediction in the vegetation cover that protects rodents from natural predation Reduction in the vegetation cover that protects rodents from natural predation Reduction of natural predation by rodents through the use of fences restricting access to plots of land Biological regulation of plant pathogenic nematodes by other nematode species Consumption of lanets, larvae or molluscs Consumption of infested fruit that has fallen to the ground Optimize as a result of reduced grass cover Appearance of insects by rubbing against trunks Consumption of auxiliary species (unknown mechanisms) Consumption of auxiliary species (unknown mechanisms) Consumption of indexted nurbed dealeases or fruit that has fallen to the ground Readuction in dualidiary species (unknown mechanisms) Consumption of auxiliary species (unknown mechanisms) Reduction of auxiliary species (unknown mechanisms) Consumption of infected print that has fallen to the ground Reduction of auxiliary species due to changes in flora Reduction of incoulum by the degradation of deal leaves or fruit that has fallen to the ground thanks to urine (no flore to inconcluste by removing low branches Reduction of inocculum by the degradation of deal leaves or fruit that has fallen to the ground thanks to urine (no flore this of ascospore projection due to removal of low branches Reduction of inocculum by the degradation of deal leaves of the weak from the vine shoulds Reduction functionate by removing the leaves from the vine shouts Reduction di nemicolimate by removing the leaves from the vine shouts Re		- Destruction of galleries by trampling	5		1						(?) 2		
- Rodent consumption - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -		- Destroying galleries by excavating the ground									(?) 2		
Pests - Reduction in the vegetation cover that protects rodents from natural predation 2 1 I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I		- Rodent consumption									(?) 2		
Pests - Reduction of natural predation by rodents through the use of fences restricting access to plots of land - 1 - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - </td <td></td> <td>- Reduction in the vegetation cover that protects rodents from natural predation</td> <td>2</td> <td></td> <td>1</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>(?) 1</td> <td></td> <td></td>		- Reduction in the vegetation cover that protects rodents from natural predation	2		1						(?) 1		
Pests - Biological regulation of plant pathogenic nematodes by other nematode species 1 Image: Consumption of insects, larvae or molluscs Image: Consumptin of insects, larvae or molluscs <t< td=""><td></td><td>- Reduction of natural predation by rodents through the use of fences restricting access to plots of land</td><td></td><td></td><td></td><td>1</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></t<>		- Reduction of natural predation by rodents through the use of fences restricting access to plots of land				1							
- Consumption of insects, larvae or molluscs - 2 (?) 13 4 4 2 (?) 1 1 - Consumption of infested fruit that has fallen to the ground (?) 9 2 (?) 8 (?) 6 (?) 2 (?) 1 1 (?) 4 - Disturbance of insects by rubbing against trunks 2 - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -	Pests	- Biological regulation of plant pathogenic nematodes by other nematode species	1										
- Consumption of infested fruit that has fallen to the ground (?) 9 2 (?) 8 (?) 6 (?) 2 (?) 1 1 (?) 4 - Disturbance of insects by rubbing against trunks 2 1 1 1 1 (?) 4 1 (?) 4 1 (?) 4 1 (?) 4 1 (?) 4 1 (?) 4 1 (?) 4 1 (?) 4 1 (?) 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		- Consumption of insects, larvae or molluscs			2	(?) 13	4	4		2	(?) 1	1	
- Disturbance of insects by rubbing against trunks 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		- Consumption of infested fruit that has fallen to the ground	(?) 9		2	(?) 8	(?) 6	(?) 2	(?) 1	1	(?) 4		
- Disturbance to insect habitat and microclimate as a result of reduced grass cover 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		- Disturbance of insects by rubbing against trunks	2										
- Appearance of insect pests living in animal dung 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		- Disturbance to insect habitat and microclimate as a result of reduced grass cover				1	1	1				1	
Auxiliary species 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	τ	- Appearance of insect pests living in animal dung			1								
Auxiliary species - Consumption of auxiliary speci		- Reappearance of auxiliary species (unknown mechanisms)	1										
species - Changes in auxiliary species due to changes in flora (?) 1 Image: Comparison of the changes in flora (?) 1 Image: Comparison of the changes in flora (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 4 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?	Auxiliary	- Consumption of auxiliary species				(?) 2							
- Reducing inoculum by trampling or eating dead leaves or fruit that has fallen to the ground (?) 9 (?) 3 (?) 2 (?) 4 (?) 4 (?) 4 - Reduction of inoculum by the degradation of dead leaves that have fallen to the ground thanks to urine (droppings in the case of poultry). (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 <td>species</td> <td>- Changes in auxiliary species due to changes in flora</td> <td>(?) 1</td> <td></td>	species	- Changes in auxiliary species due to changes in flora	(?) 1										
Pathogen - Reduction of inoculum by the degradation of dead leaves that have fallen to the ground thanks to urine (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1 (?) 1	Pathogens	- Reducing inoculum by trampling or eating dead leaves or fruit that has fallen to the ground	(?) 9			(?) 3	(?) 2				(?) 4	(?) 2	
- Improving the microclimate by removing low branches 5		- Reduction of inoculum by the degradation of dead leaves that have fallen to the ground thanks to urine (droppings in the case of poultry).	(?) 1			(?) 1	(?) 1				(?) 1	(?) 1	
- Reduced risk of ascospore projection due to removal of low branches 2 Improving the microclimate by removing the leaves from the vine shoots 1 Improving the microclimate by removing the leaves from the vine shoots 1 Improving the microclimate by removing the leaves from the vine shoots 1 Improving the microclimate by removing the leaves from the vine shoots 1 Improving the microclimate by removing the leaves from the vine shoots 1 Improving the microclimate by removing the surface 1 <td>l'unogenia</td> <td>- Improving the microclimate by removing low branches</td> <td>5</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td> </td>	l'unogenia	- Improving the microclimate by removing low branches	5										
- Improving the microclimate by removing the leaves from the vine shoots 1 I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I I		- Reduced risk of ascospore projection due to removal of low branches	2										
- Root damage caused by "ploughing" the soil or scratching the surface (?) 3 1 7		- Improving the microclimate by removing the leaves from the vine shoots	1										
		- Root damage caused by "ploughing" the soil or scratching the surface				(?) 3		1			7		
- Degradation of the tree by barking or pecking 2 17 3 2 (?) 6 3 3 1 2 2		- Degradation of the tree by barking or pecking	17	3	2	(?) 6	3	3	1	2		2	
- Damage to low branches caused by consumption, rubbing or leaning against the tree 18 5 4 3 3 2 1 2		- Damage to low branches caused by consumption, rubbing or leaning against the tree	18	5	4	3	3	2	1	2			
- Consumption of buds, leaves or fruit 20 5 4 (?) 7 3 2 1 2 2		- Consumption of buds, leaves or fruit	20	5	4	(?) 7	3	2	1	2	2		1
- Damage to the trunk by pressure or friction 4 2 (?) 4 2 1 1 1 1		- Damage to the trunk by pressure or friction	4		2	(?) 4	2	1	1	1			
Dials of domagoe - Damage to the tree by perching 2 1	Dick of domogoa	- Damage to the tree by perching				2		1					
Risk of damages 1 1 1 - Damage to vine shoots 1 1 1	RISK OF damages	- Damage to vine shoots	1	1							1		
- Soil degradation through compaction or the creation of ruts 13 2 4 2 1 1 7	- and the second se	- Soil degradation through compaction or the creation of ruts	13	2	4	2	1	1			7		
- Appearance of bare soil 2 (?) 4		- Appearance of bare soil	2			(?) 4							
- Damage to fruit 1 (?) 2 (?) 1		- Damage to fruit	1			(?) 2		(?) 1					
- Damage to young plants 5 1 1 3 2 2 1 1		- Damage to young plants	5	1	1	3	2	2	1	1			
- Degradation of trees through local over-fertilisation 2	1	- Degradation of trees through local over-fertilisation	2										

Major knowledge gaps

Animal species that have not been studied much, yet seem interesting

- Poultry: some knowledge of hens, chickens and geese, but very little is known about other species (guinea fowl, ducks, turkeys).
- Rabbits: could be an avenue, but what about outlets in the industry
- Equine ?

Mechanisms that are still poorly studied or understood

- Little is known about the impact on diseases, which is difficult to assess and quantify. Yet farmers are asking real questions on this subject.
- Impact on auxiliary species is unknown



Conclusion

- Many fruit and wine growers are motivated to introduce animals, but are concerned about damage on trees and soil
- Introducing animals can be a solution for managing weed, pest and diseases and lead to reduction of chemical treatments
- However, not all species can be equally used and mechanisms of pest and disease control remain largely unknown
- The use of animals to maintain perennial (or annual) crops requires skills in animal management and raises a number of organizational issues between livestock farmers and crop producers



Thank you for your attention



You are welcome to visit the stand of GIS Avenir Elevages (N°9)



∂venir *elevages*















