


The effect of a phytogenic on broiler performance and feed costs in normal and reduced energy diets

Marlien de Kock¹, Marçal Verdú², Sandra van Kuijk¹, Xavier Soldevila² and Georgina Farré²

¹Selko, Amersfoort, The Netherlands. ²BonÀrea Agrupa, Animal Nutrition and Feed Industry, Guissona, Spain.







Plant metabolites

Primary compounds:
Directly involved in growth,
development and
reproduction

Secondary compounds: Help the plant to survive in its environment; broad functional spectrum.







Inutreco

The gut is a complex sensory organ, with a broad range of receptors capable of responding to plant secondary metabolites

Using this concept, a formulated blend of essential oils (Fytera® Perform) improved broiler performance in field-like conditions



Science-based, proven mode of action










Fytera Perform reduced expression of proinflammatory genes and increased expression of genes related to gut barrier function




Lillehoj (USDA), Wall et al., unpublished results; Doses used in vitro were equivalent to those that would be fed in vivo

Meta-analysis of performance data



Fytera Perform improves FCR and Body Weight



Fytera Perform (25 g/t) improved on average

- FCR by -5 points
- Live BW by +45 g

Dietary copper levels did not affect product efficacy.

Trial detail



<u>Research Collaboration</u> Corporación Alimentaria Guissona, S.A. - bonÀrea Agrupa.



Andorra

bonArea - Corporacio
Alimentaria (Guissona

o Terrassa

o Barcelona

Reus o Tarragona

Trial Site: Guissona, Spain

Objective: Validate growth performance of Fytera Perform in a commercial or cost-saving diet

KPI: Growth performance, carcass performance and quality, economics

Trial design





3,072 Ross 308



36 days



wheat-soybean-corn based diet



3 treatments - 8 pens/treatment - 128 birds/pen

- T1: Control (basal diet)
- T2: FP (basal diet + 25 g/t Fytera Perform)
- T3: FP-dil (diluted basal diet + 25 g/t Fytera Perform)

Trial detail



Standard feed formulation

Feed Phase	Energy (kcal/kg)	Protein (%)
Pre-starter	3 001	21.7
Starter	3 030	19.2
Grower	3 048	18.5
Finisher	3 081	17.0


Diluted diets

Nutrient	0-7 d	8-21 d	21-30 d	31-36 d
AME, Kcal	-15	-20	-25	-30
dLys*	=	=	-2%	-2%
Ca total	0.95	0.75	0.70	0.65
P dig	0.46	0.40	0.36	0.31

^{*} Rest of AA corrected according to dLys


Mortality






Performance






Performance






Performance – Feed Conversion Ratio





Carcass weight






 There were numerical differences in carcass weight, close to tendency (P = 0.12)

Economics



Fytera Perform Cost = 0,007 Euro



Conclusion



- Both T2 and T3 improved growth performance compared to the control diet (T1)
- T3 was the best treatment concerning cost-effectiveness when compared to the control diet (T1) and T2

Hypothesis.

"Phytogenics can improve gut health in such a way as to improve nutrient uptake and reduce nutrient requirements in feed to enable producers to grow chickens more economically"



Thank you



