










## The planet is heating up...










## Impact of climate change in dairy cows



biorender.com

\*5-20% in following lactations


- Heat stress (HS) negatively affects production and health of dairy cows.
- In the U.S., HS costs \$1.5B/year
- In Europe, -5.4% the monthly income in the summer season







## Threshold of heat stress: temperature humidity index (THI) > 68

















### Rumination can serve as an early indicator of heat stress





 A HS threshold of THI 52 was identified based on rumination time using broken-stick regression, but lacking flexibility.







# Explore the effect of THI on rumination in comparison to milk production using a more flexible modeling framework

### Hypothesis:

- the HS threshold of rumination time is lower than that of milk production,
- the HS threshold can vary depending on
  - Parity
  - Production level of animals
  - Lactation stage

Item	Pa	rity	Production level		
	Primiparous	Multiparous	Low	High	
Number of cows, n	•	•		Ü	
Rumination time	498	546	99	117	
Milk yield	474	495	99	117	
Milk quality measures <sup>2</sup>	696	605	66	88	
Mean (±SD)					
Parity	1.0 (±0.00)	2.2 (±0.42)	2.33 (±0.47)	2.24 (±0.42)	
Rumination time, min/d	542 (±69.8)	524 (±78.4)	512 (±76.9)	547 (±72.3)	
Milk yield, kg/d	35.4 (±6.34)	40.3 (±10.32)	35.8 (±7.30)	51.2 (±6.20)	







#### **Data collection**

 Data were collected from 920 lactating dairy cows (Holsteins) for 1,108 days between May 2015 to June 2018 in Beijing, China.

SCR HR sensors were used

### Exploration of different climate metrics

- Daily mean THI
- Number of hour per day when THI > 68
- Daily max or min THI

Climate metrics	Observations, n	Mean	Median	Min	Max	SD
THI	1,108	56.4	59.2	25.6	84.8	15.46
THI above 68, h/d	1,108	7.18	0.00	0.00	24.0	8.996









# Generalized additive models (GAMs) – a semi-parametric approach

GAMs are smooth semi-parametric models of the form:

$$g(\mathbb{E}[y|X]) = \beta_0 + f_1(X_1) + f_2(X_2, X_3) + \dots + f_M(X_N)$$

- GAMs extend generalized linear models by allowing non-linear functions of features
- Since the model is additive, it is easy to examine the marginal effect of each variable










## Model structure and determination of inflection point

- Generalized additive mixed model including:
  - Random effect of cows
  - Fixed effect of days in milk (interacting with parity or production level)
  - Fixed effect of THI (interacting with parity or production level)
  - Fixed effect of time of year/Julian day (but dropped due to lack of effect)




- Pointwise derivatives (i.e., the slopes) of the fitted means determined based on the partial effects of THI.
- Inflection points will then be determined from the slopes and will be used for interpreting the HS threshold for different parameters.








## Variable heat stress thresholds for primi- vs. multiparous cows








## Variable heat stress thresholds for high- vs. low-producing cows







### Would a different climate metrics tell us more?







## Take home message

- Rumination can be considered as an early indicator of heat stress.
- Heat stress threshold based on rumination time showed THI = 58.8 regardless of parity of cows, but limited to up to ~2 lactations; compared to THI = 65 for multiparous cows.
- Heat stress threshold based on rumination time showed that high-producing cows are more sensitive than low-producing cows (THI = 60.9 vs. 58.6, respectively).
- Further investigation and tuning to improve model robustness.









Yachun Wang



Aira Serviento



**Daniel Rico** 

**Environment** 

## Thank you!



Contact: Mutian Niu mutian.niu@usys.ethz.ch