Performances of primiparous rabbit does subjected of dietary restriction during gestation

Victor Pinheiro. José Teixeira. Divanildo Monteiro

UTAD/CECAV - Vila Real. Portugal











Introduction

The ad libitum feeding of rabbit does, especially in the first production cycle can lead to obesity and may cause reproductive problems and increasing mortality in the first parturition.

Dietary restriction during certain periods of does development can be advantageous. Reducing this problem and saving on feeding costs.

Objective

Dietary restriction was applied to two groups of nulliparous does between the 4th and 25th day of gestation.

Material and methods



Animals:

- 60 nulliparous rabbits does with 18 weeks (NZ X C)
- 2 groups (30 animals/group)

2 Treatments with a dietary restriction (25%)

- **R7D** feed restriction every days (136g/d)
- R6F feed restriction in 6 days (159 g/day) and 1 day of fasting

From 4th to 25th day of gestation. Other days - feed ad libitum

Material and methods





Measurements:

Litter weight:

at birth,11 and 20 days postpartum and at weaning

Rabbit does weight:


at partum, 11 and 20 days postpartum and at weaning

Were monitored:

Rabbit does fertility and the daily mortality of pups







In R6F group

At partum the fertility significantly (P<0.05) increased from 86% to 100%.

Results – Live weight of rabbits and litters

	Treatment		P
	R7D	R6F	
Litter Weight			
At birth	577.3	583.5	ns
Day 1	592.0	594.4	ns
Day 11	1823.7	1724.9	ns
Day 20	2944.7	2853.7	ns
Weaning (34d)	7665.4	7716.4	ns
Rabbit weight			
Day 1	53.8	54.0	ns
Day 11	172.8	168.2	ns
Day 20	285.4	285.0	ns
Weaning (34d)	747.2	788.9	0.045

In the Rabbit R6F group

Live weight at weaning (34 d of age) was significantly (P=0.045) higher (more 41g)

Results – number of rabbits

	Treatment		D.
	R7D	R6F	P
Live birth	11.0	11.5	ns
Death birth	0.64	1.43	ns
Total birth	11.6	13.0	0.07
Day 1	11.0	11.0	ns
Day 11	10.6	10.3	ns
Day 20	10.3	10.1	ns
Weaning (34d)	10.2	9.9	ns

In the Rabbit **R6F** group

Prolificity increased (P=0.07), from 11.6 to 13.0 total births

Introduction	Methodology	Results	Conclusion	Considerations
--------------	-------------	---------	------------	----------------


Results – gestation time

	Treatment		
	R7D	R6F	Р
Gestation time (d)	30.1	29.9	ns
Gestation time (h)	726.6	719.9	0.052

In the R6F

Gestation time showed a tendency (P=0.052) to decrease approximately 7 hours





Mortality of the pups during the suckling period did not differ and remaining around 8.5%, but it's 3.7% higher in the group R6F.



Implementing a fasting day during the gestation period in nulliparous does improves fertility and not change the others parameters.

This practice can make rabbit farming easier.



Considerations

In future work

- complement with other measurements: milk, blood ...
- productive longevity
- productivity in the following cycles





Financing : Project UIDB/ 00772/2020 e LA/P/0059/2020 financed by Fundação Portuguesa para a Ciência e Tecnologia (FCT).