

Implementing an autodrafter-weighing scale for assessing the body weight changes and grouping dairy sheep and goats during lactation

G. Caja¹, A. Elhadi¹ and R. Costa²

¹Group of Research in Ruminants, Department of Animal and Food Sciences

²Servei de Granges i Camps Experimentals

Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain







TechCare: Integrating innovative **TECH**nologies along the value **C**hain to improve small ruminant welf**ARE** management UE H2020 Project Contract No 862050: www.techcare-project.eu



 Prioritizations in dairy sheep and goats: Intensive conditions in Spain (Caja & Elhadi, 2021, 2022)

| Dairy sheep/goats | Priority | Votes |
|--|----------|-------|
| Welfare issues | | |
| Mastitis | 1 | 79% |
| Nutrition issues | 2 | 69% |
| Environment and shelter conditions | 3 | 69% |
| Sensors and technologies | | |
| Weather stations internal-external (THI) | 1 | 83% |
| Electronic milk meters (milk yield) | 2 | 68% |
| Automated weighing scales (BW) | 3 | 56% |
| Acelerometrers-3A (activity) | 4 | 51% |



2

Task 3.3. Aim: Adapting and testing pre-existing PLF technologies to match specific animal welfare priorities and other production parameters


Specific aims and questions to respond:

- Commercially available device able to be shared among farms:

 Prattley 2-ways Autodrafter Datamars (12,000 €, software and TVA included)
- Able to be implemented in previously existing farms:
 SGCE of UAB (Bellaterra)
- Able to be automatically operated in dairy ewes and dairy goats:
 Do not alter operation sheep and goat flow after milking (≈200 head/h?)
- User friendly software, robustness and durability:
 On place from September 2023 (12 mo currently at the UAB)
- Use of weekly BW as welfare indicator: Nutrition/diseases issues (EWS?)
- Able to be used for other welfare indicators (BCS, Wool, Stress, Order,...)?




Prattley-Datamars autodrafter scale implemented at the experimental farm of the UAB




Experimental farm features and operating conditions:


- Dairy sheep (Manchega, n = 40; Lacaune, n = 120) and goats (Murciano-Granadina, n = 50)
- Semi-intensive: 6 h/d grazing + TMR (F:C = 60:40) + 0.1 kg concentrate at each milking.
- Joined as an only flock for grazing (6 h/d) and milking (2x daily).
- Weighed once a week after milking and before grazing during the whole year.
- Groups recomposed after grazing.















•

Results of the Autodrafter-weighing throughput






Learning progress by sessions






Upload into the Datamars Livestock cloud (free)



Individual analysis of tails: 1/2



TechCare

Individual analysis of tails: 2/2



Conclusions:

1) Prattley 2-ways Autodrafter- Datamars:

- Was easily implemented in a preexisting farm for sheep and goats, despite the insufficient instructions received (to be improved!).
- The use of EU official LF-ISO readers (134.2 kHz) and eID boluses (22 g) were fully efficient for automatic weighing and sorting of dairy sheep and goats.
- Both ewes and does were easily trained and reached a throughput >300 head/h after 15 sessions.
- Differences by species, breed (and individuals?)

2) Datamars Livestock software:

• Was easily adapted as an EWS to detect ewes and does loosing weight (or winning excessive weight) during lactation.



Thank you for attention!

gerardo.caja@uab.es



www.techcare-project.eu



Project funded by the H2020 research and innovation program of the European Union (Contract No 862050)

